

Tuesday, Feb. 13th

**NO SCHOOL ON MONDAY,
FEB. 19TH FOR FAMILY DAY**

Plan For Today:


1. Any questions from Ch3?

- * Hand-in Ch3 Project
- * Do Chapter 3 Test

2. Start Chapter 4: Rational Functions

- * 4.3: Rational Functions
- * 4.4: Graphing Rational Functions

3. Work on Practice Questions from Workbook



START UNIT 1 REWRITE AT ~ 12:10PM

Plan Going Forward:

1. Finish going through all of the 4.3 questions in workbook and start working on practice review handout.

● **CHECK-IN QUIZ ON 4.3 ON THURSDAY, FEB. 15TH**

2. We will finish Ch4 on Thursday (hopefully).

- **CHAPTER 4 PROJECT DUE TUESDAY, FEB. 20TH**
- **CHAPTER 4 TEST ON TUESDAY, FEB. 20TH**

* **UNIT 2 EXAM ON CH3&4 ON THURSDAY, FEB. 22ND**

- 10 Multiple Choice & 20 marks on the Written
- ~1 hour - please prepare so you are not "learning" while doing the test
- Closed-book - no notes
- Rewrite is following Tuesday after class at 12:30pm
- I will email you this weekend when marks are posted so you can decide on the rewrite
- I will go over the marked exam on Tuesday

Please let me know if you have any questions or concerns about your progress in this course. The notes from today will be posted at anurita.weebly.com after class.
Anurita Dhiman = adhiman@sd35.bc.ca

Tuesday, Feb. 13th In-Class Notes



Sec9.1 Intro
Mar2023

43


Section 2.1: Intro to Graphing Rational Functions with Transformations

Graph the following functions on the grid provided. Show your base function table of values and the transformed function's table of values. Also include the domain and range.

1. $y = \frac{1}{x}$ This is the base function of a rational function before transformations

x	y
-3	$-\frac{1}{3}$
-2	$-\frac{1}{2}$
-1	-1
$-\frac{1}{2}$	-2
0	\emptyset
$\frac{1}{2}$	2
1	1
2	$\frac{1}{2}$
3	$\frac{1}{3}$

$y = \frac{1}{-\frac{1}{2}}$
 $= -2$
 $y = \frac{1}{0} = \emptyset$
 undefined
 = asymptote



Domain: $\{x \mid x \neq 0, x \in \mathbb{R}\}$

Range: $\{y \mid y \neq 0, y \in \mathbb{R}\}$.

Equations of the asymptotes: horizontal asymptote (HA) $\rightarrow y=0$
 vertical asymptote (VA) $\rightarrow x=0$

This is the standard form of a rational function for graphing transformations:

Base Function $y = \frac{1}{x} \rightarrow y = \frac{a}{b(x-h)} + k$ $(\frac{a}{b}) = V$ stretch

h and k determine the location of the asymptotes.

$$2. y = \frac{3}{x-4} + 2$$


Transformations:

- ① VE of 3
- ② 4 right, 2 up.

Base:

x	$y = \frac{1}{x}$	$3y$	$x+4$	$3y+2$
-3	$-\frac{1}{3}$	-1	1	1
-2	$-\frac{1}{2}$	$-\frac{3}{2}$	2	$\frac{1}{2}$
-1	-1	-3	3	-1
$-\frac{1}{2}$	-2	-6	$\frac{7}{2}$	-4
0	\emptyset	\emptyset	4	\emptyset
$\frac{1}{2}$	2	6	$\frac{9}{2}$	8
1	1	3	5	5
2	$\frac{1}{2}$	$\frac{3}{2}$	6	$\frac{7}{2}$
3	$\frac{1}{3}$	1	7	3

$$\frac{1}{2} + \frac{8}{2}$$



Domain: $\{x \mid x \neq 4, x \in \mathbb{R}\}$.

Range: $\{y \mid y \neq 2, y \in \mathbb{R}\}$.

Asymptotes: VA: $x=4$ HA: $y=2$

never write

~~VA=4 HA=2~~

Simplify the following by factoring and state the non-permissible values (NPVs or Restrictions):

1. $y = \frac{x^2 - 3x}{x}$

- ① Factor
- ② Simplify
- ③ NPV from original b/f simplifying

① $y = \frac{x(x-3)}{x}$

② $y = x - 3$

③ NPV $x \neq 0$

2. $y = \frac{x+4}{3x^2 + 10x - 8}$

① $y = \frac{x+4}{(x+4)(3x-2)}$

-24
+12, -2 = 10

(3x+12)(3x-2)

② $y = \frac{1}{3x-2}$

③ NPV: $x \neq -4, x \neq \frac{2}{3}$

$x+4 \neq 0$
 $\rightarrow -4$
 $x \neq -4$

$3x-2 \neq 0$
 $\rightarrow 12$
 $3x \neq \frac{2}{3}$
 $x \neq \frac{2}{9}$

3. $y = \frac{2x^2 - 4x}{x^2 + 5x - 14}$

4.3: Rational Function Transformations

Graphing the basic rational function with transformations

8-4 Rational Functions

The rational function $f(x) = \frac{1}{x}$ can be transformed by using methods similar to those used to transform other types of functions.

$|a|$ → vertical stretch or compression factor
 $a < 0$ → reflection across the x -axis

k → vertical translation
down for $k < 0$; up for $k > 0$

$$f(x) = \frac{a}{x - h} + k$$

h → horizontal translation
left for $h < 0$; right for $h > 0$

Holt Algebra 2


Copyright © by Holt, Rinehart and Winston. All Rights Reserved.

Core Concept

Graphing Translations of Simple Rational Functions

To graph a rational function of the form $y = \frac{a}{x - h} + k$, follow these steps:

- Step 1** Draw the asymptotes $x = h$ and $y = k$.
- Step 2** Plot points to the left and to the right of the vertical asymptote.
- Step 3** Draw the two branches of the hyperbola so that they pass through the plotted points and approach the asymptotes.




Rational Functions

Example 1: Transforming Rational Functions

Using the graph of $f(x) = \frac{1}{x}$ as a guide, describe the transformation and graph each function.


A. $g(x) = \frac{1}{x+2}$

Because $h = -2$,
translate f 2 units left.



B. $g(x) = \frac{1}{x} - 3$

Because $k = -3$,
translate f 3 units down.



Holt McDougal Algebra 2

Copyright © by Holt Mc Dougal. All Rights Reserved.

Rational Functions and Transformations

Graph $f(x) = \frac{5}{x-2}$.

$$f(x) = \frac{a}{x-h} + k$$


Compare the graph of $f(x) = \frac{5}{x-2}$
with $f(x) = \frac{1}{x}$.

Vertical stretch by a
factor of 5, horizontal
translation 2 units right.

Sketch the graph using transformations.
The horizontal asymptote is translated
2 units right as well.

Domain: $x \neq 2, x \in R$.

Range: $y \neq 0, y \in R$






5.6.7

1. Match the rational equation with the correct graph. Explain your rationale for how you matched the equation with the graph.

a) $y = -\frac{1}{x}$

b) $y = -\frac{1}{x-3} + 4$

c) $y = -\frac{1}{x+1} - 2$



4.3: Analysing Rational Function Graphs

Graph Rational Functions with Holes

If the degree of the numerator < degree of the denominator then **horizontal asymptote** is at $y = 0$.

The **vertical asymptotes** will occur where the denominator equals zero.


If there is a common factor in the numerator and denominator then the graph of a rational function will have a **hole** when a value of x causes both the numerator and the denominator to equal 0. We can set the common factor to zero and solve for x to find the hole.

Example:

$$f(x) = \frac{-2x + 8}{x^2 - 6x + 8}$$
$$= \frac{-2(x - 4)}{(x - 4)(x - 2)}$$

hole at $x = 4$

vertical asymptote
at $x = 2$



Sketching the Graph of a Rational Function by Hand

Guidelines for Graphing Rational Functions

1. Write the rational expression in **simplest form**, by factoring the numerator and denominator and dividing out common factors.
2. Find the coordinates of any **“holes”** in the graph.
3. Find and plot the **y-intercept**, if any, by evaluating **$f(0)$** .
4. Find and plot the **x-intercept(s)**, if any, by **finding the zeros of the numerator**.
5. Find the **vertical asymptote(s)**, if any, by **finding the zeros of the denominator**. Sketch these using dashed lines.
6. Find the **horizontal asymptote**, if any, by **comparing the degrees of the numerator and denominator**. Sketch these using dashed lines.
7. Find the **oblique asymptote**, if any, by **dividing the numerator by the denominator using long division**.
8. Plot **5-10 additional points**, including points close to each x-intercept and vertical asymptote.
9. Use **smooth curves** to complete the graph.

Rules for Graphing Rationals	Examples
<p>To get the end behavior asymptote (EBA), you want to compare the degree in the numerator to the degree in the denominator. There can be at most 1 EBA and most of the time, these are horizontal.</p> <p>➤ If the degree (largest exponent) on the bottom is greater than the degree on the top, the EBA (which is also a horizontal asymptote or HA) is $y=0$.</p>	$y = \frac{x+2}{x^2-4}$ <p>Notice that even though we can take out a removable discontinuity ($x+2$), the bottom still has a higher degree than the top, so the HA/EBA is $y=0$.</p>
<p>➤ If the degree on the top is greater than the degree on the bottom, there is no EBA/HA. However, if the degree on the top is one more than the degree on the bottom, then there is a slant (oblique) EBA asymptote, which is discussed below.</p>	$y = \frac{x^3+2}{x-4}$ <p>No HA/EBA. Vertical asymptote is still $x=4$.</p>
<p>➤ If the degree is the same on the top and the bottom, then divide coefficients of the variables with the highest degree on the top and bottom; this is the HA/EBA. You can determine this asymptote even without factoring.</p>	$y = \frac{2x^3+2}{3x^3-4}$ <p>Since the degree on the top and bottom are both 3, the HA/EBA is $y=\frac{2}{3}$.</p>
<p>➤ If the degree on the top is one more than the degree on the bottom, then the function has a slant or oblique EBA in the form $y=mx+b$. We have to use long division to find this equation.</p> <p>We can just ignore or "throw away" the remainder and just use the linear equation. Weird, huh?</p>	$y = \frac{2x^2+x+1}{x-4}$ $\begin{array}{r} 2x+9 \\ x-4 \overline{) 2x^2+x+1} \\ \underline{2x^2-8x} \\ 9x+1 \\ \underline{9x-36} \\ 37 \end{array}$ <p>EBA: $y=2x+9$</p>
<p>➤ (more Advanced) Find the point where any horizontal asymptotes cross the function by setting the function to the horizontal asymptote, and solving for "x". You already have the "y" (from the HA equation).</p>	<p>Q: Where does $y = \frac{-x^2+x}{x^2+x-12}$ intersect its EBA?</p> <p>A: Note that the EBA is $y = \frac{-1}{1} = -1$. Now set $\frac{-x^2+x}{x^2+x-12} = -1$ and cross multiply: $-x^2+x = -1(x^2+x-12)$; $x=6$.</p> <p>So the point where the function intersects the EBA is (6, -1).</p>

Rules for Graphing Rationals	Examples
<p>First factor both the numerator and denominator, and cross out any factors in both the numerator and denominator.</p> <p>➤ If any of these factors contain variables, these are removable discontinuities, or “holes” and will be little circles on the graphs. The idea is that if you cross out a polynomial, you can't forget that it was in the denominator and can't “legally” be set to 0. (We will see graph later.)</p> <p>The domain of a rational function is all real numbers, except those that make the denominator equal zero, as we saw earlier.</p> <p>(Note that if after you cross out factors, you still have that same factor on the bottom, the “hole” will turn into a vertical asymptote; follow the rules below).</p>	$y = \frac{x^2 - 5x + 6}{x - 3} = \frac{\cancel{(x-3)}(x-2)}{\cancel{(x-3)}} = x - 2$ <p>This function reduces to the line $y = x - 2$ with a removable discontinuity (a little circle on the graph) where $x = 2$ and $y = (2) - 2 = 0$ (plug 2 in for y in original or reduced fraction). So the hole is at $(2, 0)$.</p> <p>Domain is $(-\infty, 3) \cup (3, \infty)$, since a 3 would make the denominator = 0. It's like we have to “skip over” the 3 with interval notation.</p>
<p>To get vertical asymptotes or VAs:</p> <p>➤ After determining if there are any holes in the graph, factor (if necessary) what's left in the denominator and set the factors to 0. For any value of x where these factors could be 0, this creates a vertical asymptote at “$x =$” for these values.</p> <p>Note: There could a multiple number of vertical asymptotes, or no vertical asymptotes.</p> <p>Don't forget to include the factors with “x” alone ($x = 0$ is the vertical asymptote).</p>	$y = \frac{x^2 - 5x + 6}{x(x^2 - 9)} = \frac{\cancel{(x-3)}(x-2)}{x\cancel{(x-3)}(x+3)} = \frac{x-2}{x(x+3)}$ <p>Vertical asymptotes occur when $(x-0) = 0$ or $(x+3) = 0$, or $x = 0$ or $x = -3$.</p> <p>Domain is $(-\infty, -3) \cup (-3, 0) \cup (0, 3) \cup (3, \infty)$, since anything that could make the denominator 0 (even a hole) can't be included. So we have to “skip over” $-3, 0,$ and 3.</p>

Steps		Graph
$y = \frac{3x^2 - 8x - 16}{3x^2 - 16x + 16}$ <p>Factor:</p> $y = \frac{(3x+4)(\cancel{x-4})}{(3x-4)(\cancel{x-4})}$ <p>Removable Discontinuity or Hole: $x = 4$, plug in 4 for x to get $y = 2$. RD is $(4, 2)$.</p> <p>VA: Set denominator to 0 after removing the hole; $3x - 4 = 0$ $x = \frac{4}{3}$.</p>	<p>EBA/HA: Since the degree is the same on the top and bottom (both are 2), we take the coefficients and divide them: $y = \frac{3}{3} = 1$.</p> <p>x-intercept (root): Set y (or top) to 0: $3x + 4 = 0$; $x = -\frac{4}{3}$ $(-\frac{4}{3}, 0)$</p> <p>y-Intercept: Set x to 0: $y = \frac{3(0) + 4}{3(0) - 4} = -1$ $(0, -1)$</p> <p>Domain: Can't be any value of x that makes the bottom zero: $(-\infty, \frac{4}{3}) \cup (\frac{4}{3}, 4) \cup (4, \infty)$</p>	<p>$y = \frac{3x^2 - 8x - 16}{3x^2 - 16x + 16}$</p> <p>Graph showing the function $y = \frac{3x^2 - 8x - 16}{3x^2 - 16x + 16}$. The graph has a vertical asymptote at $x = \frac{4}{3}$ and a horizontal asymptote at $y = 1$. The x-intercept is $(-\frac{4}{3}, 0)$ and the y-intercept is $(0, -1)$. There is a removable discontinuity (hole) at $(4, 2)$.</p>
$y = \frac{x}{x^2 + 4x - 5}$ <p>Factor:</p> $y = \frac{x}{(x+5)(x-1)}$ <p>VA: Set denominator to 0 after factoring; we have 2 of them: $x = -5$, $x = 1$.</p> <p>EBA/HA: Since the degree on the top (1) is less than the degree on the bottom (2), the EBA or VA is $y = 0$.</p>	<p>x-intercept (root): Set y (or top) to 0: $x = 0$. $(0, 0)$</p> <p>y-Intercept: Set x to 0: $y = \frac{(0)}{(0)^2 + 4(0) - 5} = 0$: $(0, 0)$</p> <p>"T-chart": Try some points around the vertical asymptotes: $x = -6$, $y = -.86$ $x = -4$, $y = .8$ $x = 2$, $y = .29$</p> <p>Domain: Can't be any value of x that makes the bottom zero: $(-\infty, -5) \cup (-5, 1) \cup (1, \infty)$.</p>	<p>$y = \frac{x}{x^2 + 4x - 5}$</p> <p>Graph showing the function $y = \frac{x}{x^2 + 4x - 5}$. The graph has vertical asymptotes at $x = -5$ and $x = 1$, and a horizontal asymptote at $y = 0$. The x and y intercept is $(0, 0)$. Other points plotted include $(-6, -.86)$, $(-4, .8)$, and $(2, .29)$.</p>


Rules for Asymptotes

Exponent of highest degree term in denominator larger than highest degree term in numerator then Horizontal Asymptote is

$$y = 0$$


Exponent values of highest degree of terms in numerator and denominator the same then Horizontal Asymptote is

$$y = \text{ratio of their coefficients}$$



2.2 Point of Discontinuity

$$f(x) = \frac{x^2 - x - 2}{x - 2}$$







5. Which graph matches each rational function? Explain your choices.

a) $A(x) = \frac{x^2 + 2x}{x^2 + 4}$

b) $B(x) = \frac{x - 2}{x^2 - 2x}$

c) $C(x) = \frac{x + 2}{x^2 - 4}$


d) $D(x) = \frac{2x}{x^2 + 2x}$



8. Write the equation of a possible rational function with each set of characteristics.

- a) vertical asymptotes at $x = \pm 5$ and x-intercepts of -10 and 4
- b) a vertical asymptote at $x = -4$, a point of discontinuity at $\left(-\frac{11}{2}, 9\right)$, and an x-intercept of 8
- c) a point of discontinuity at $\left(-2, \frac{1}{5}\right)$, a vertical asymptote at $x = 3$, and an x-intercept of -1
- d) vertical asymptotes at $x = 3$ and $x = \frac{6}{7}$, and x-intercepts of $-\frac{1}{4}$ and 0

Graphing Practice



Characteristics of Rational Functions Worksheet

1. $f(x) = \frac{x^2 + x - 2}{x^2 - x - 6}$


① Factor $f(x) = \frac{(x+2)(x-1)}{(x-3)(x+2)}$

② Simplify $f(x) = \frac{x-1}{x-3}$

③ Characteristics.


Characteristic	Answer
a. simplified function	$f(x) = \frac{x-1}{x-3}$
p.179 b. POD (HAE) Point of Discontinuity	$(x+2) \rightarrow x \neq -2$ $(-2, \frac{3}{5})$
p.176 c. vertical asymptote(s)	$(x-3)$ VA $\rightarrow x=3$
p.176 #2 d. horizontal asymptote	$\frac{1x^2}{1x^2} \rightarrow y=1$ HA
p.178 e. y-intercept	$x=0$ $y = \frac{0-1}{0-3} \rightarrow y = \frac{1}{3}$ $(0, \frac{1}{3})$
p.178 f. x-intercept (numerator)	$y=0 \rightarrow 0 = \frac{x-1}{x-3}$ $0 = x-1 \rightarrow x=1$ $(1, 0)$
g. domain VA + POD	$\{x \mid x \neq 3, x \neq -2, x \in \mathbb{R}\}$
h. range HA + POD	$\{y \mid y \neq 1, y \neq \frac{3}{5}, y \in \mathbb{R}\}$

$y = \frac{-2-1}{-2-3}$
 $y = \frac{-3}{-5}$
 $y = \frac{3}{5}$
when $x=-2$




2. $f(x) = \frac{2x^2}{x^2 - 1}$

Characteristic	Answer
a. simplified function	
b. POD	
c. vertical asymptote(s)	
d. horizontal asymptote	
e. y-intercept	
f. x-intercept	
g. domain	
h. range	




3. $f(x) = \frac{3}{x-2}$

Characteristic	Answer
a. simplified function	
b. POD	
c. vertical asymptote(s)	
d. horizontal asymptote	
e. y-intercept	
f. x-intercept	
g. domain	
h. range	




4. $f(x) = \frac{2x-1}{x}$

Characteristic	Answer
a. simplified function	
b. POD	
c. vertical asymptote(s)	
d. horizontal asymptote	
e. y-intercept	
f. x-intercept	
g. domain	
h. range	




5. $f(x) = \frac{x^2 + x - 12}{x^2 - 9}$

Characteristic	Answer
a. simplified function	
b. POD	
c. vertical asymptote(s)	
d. horizontal asymptote	
e. y-intercept	
f. x-intercept	
g. domain	
h. range	




6. $f(x) = \frac{x^2 - 4}{x + 3}$

Characteristic	Answer
a. simplified function	
b. POD	
c. vertical asymptote(s)	
d. horizontal asymptote	
e. y-intercept	
f. x-intercept	
g. domain	
h. range	




$$7. f(x) = \frac{x^2 - x}{x + 1}$$

Characteristic	Answer
a. simplified function	
b. POD	
c. vertical asymptote(s)	
d. horizontal asymptote	
e. y-intercept	
f. x-intercept	
g. domain	
h. range	




8. $f(x) = \frac{x^2 - x - 2}{x - 1}$

Characteristic	Answer
a. simplified function	
b. POD	
c. vertical asymptote(s)	
d. horizontal asymptote	
e. y-intercept	
f. x-intercept	
g. domain	
h. range	



9. $f(x) = \frac{x+1}{x^2+3x+2}$

Characteristic	Answer
a. simplified function	
b. POD	
c. vertical asymptote(s)	
d. horizontal asymptote	
e. y-intercept	
f. x-intercept	
g. domain	
h. range	



10. $f(x) = \frac{x^2 - 9}{x^2 - 2x - 3}$

Characteristic	Answer
a. simplified function	
b. POD	
c. vertical asymptote(s)	
d. horizontal asymptote	
e. y-intercept	
f. x-intercept	
g. domain	
h. range	

